

ANNEXE: 1

Prevalence of active syphilis over six rounds of serological surveillance, 1998-1999, 1999-2000, 2000-2001, 2002, 2003-2004, 2004-2005 and 2006

Study Populations, Geographical Location	Active syphilis % Positive (95% CI), number positive (total number tested)						
	1998-1999 Round I	1999-2000 Round II	2000-2001 Round III	2002 Round IV	2003-2004 Round V	2004-2005 Round VI	2006 Round VII
Injection Drug Users (IDU):							
Detoxification Clinic:							
Central-A	4.5 (2.7-7.0), 18 (402)	4.0 (2.3-6.4), 18 (402)	4.3 (1.2-10.8), 4 (92)	ND	ND	ND	ND
Out of detoxification clinic:							
Central- A [†]	ND	9.3 (6.7-12.5), 39 (418)	2.5 (1.2-4.5), 10 (401)	3.5 (1.9-5.8), 14 (403)	1.2 (0.4-2.9), 5 (404)	2.9 (2.0-4.1), 31 (1061)	2.3 (1.5-3.4), 25 (1066)
Central-A1	ND	ND	ND	ND	ND	3.8 (2.5-5.5), 25 (664)	3.7 (2.4-5.5), 25 (668)
Central-A2	ND	ND	ND	ND	ND	1.5 (0.6-3.3), 6 (397)	0 (398)
Central-C	ND	ND	ND	ND	ND	1.8 (0.7-3.6), 7 (395)	0.7 (0.1-2.4), 2 (301)
Central-E	ND	ND	ND	ND	5.6 (2.1-11.8), 6 (107)	1.0 (0-5.3), 1 (103)	2.9 (0.6-8.1), 3 (105)
Central-H	ND	ND	ND	ND	1.6 (0.2-5.8), 2 (122)	3.9 (1.6-7.9), 7 (178)	4.4 (1.8-8.8), 7 (160)
Southeast-D	ND	ND	ND	ND	7.0 (2.6-14.6), 6 (86)	2.5 (0.7-6.3), 4 (159)	2.8 (0.9-6.5), 5 (177)
Southeast-H1	ND	ND	ND	ND	ND	9.0 (5.0-14.7), 14 (155)	5.8 (2.4-11.6), 7 (120)
Northwest-A	ND	4.1 (2.4-6.5), 17 (416)	1.5 (0.5-3.2), 6 (402)	1.7 (0.7-3.5), 7 (405)	1.3 (0.4-2.9), 5 (394)	1.0 (0.3-2.3), 4 (398)	1.3 (0.4-2.9), 5 (393)
Northwest-B	ND	ND	1.7 (0.2-5.9), 2 (120)	2.0 (0.5-5.0), 4 (200)	1.7 (0.5-4.2), 4 (239)	0.5 (0-2.6), 1 (208)	0.5 (0-2.8), 1 (200)
Northwest-B1	ND	ND	ND	ND	1.3 (0-6.9), 1 (78)	ND	ND
Northwest-B2	ND	ND	ND	ND	2.1 (0.1-11.3), 1 (47)	1.5 (0-8.2), 1 (66)	4.3 (0.9-12.2), 3 (69)
Northwest-C	ND	ND	ND	ND	ND	ND	2.7 (0.9-6.1), 5 (187)
Northwest-D	ND	ND	ND	ND	ND	0 (120)	0.5 (0-2.9), 1 (193)
Northwest-F	ND	ND	ND	ND	0 (85)	0 (57)	1.4 (0-7.8), 1 (69)
Northwest-F1	ND	ND	ND	ND	3.5 (0.4-12.1), 2 (57)	0 (49)	1.8 (0-9.7), 1 (55)
Northwest-G	ND	ND	ND	ND	ND	1.8 (0.2-6.4), 2 (111)	0.8 (0-4.5), 1 (122)
Northwest-K	ND	ND	ND	ND	ND	ND	1.1 (0.2-3.1), 3 (279)
Southwest-B	ND	ND	ND	ND	ND	3.0 (0.6-8.5), 3 (100)	2.3 (0.5-6.5), 3 (132)
Southwest-D	ND	ND	ND	ND	ND	1.0 (0.1-3.5), 2 (201)	0.9 (0.1-3.2), 2 (226)
South-A	ND	ND	ND	ND	ND	0 (202)	0 (234)
Female IDU:							
Central-A, E, H	ND	ND	ND	ND	ND	9.2 (4.7-15.9), 11 (119)	9.9 (5.2-16.7), 12 (121)
Heroin Smokers:							
Central-A	ND	ND	ND	3.4 (1.8-5.7), 13 (387)	2.6 (1.2-4.7), 10 (391)	3.0 (1.6-5.2), 12 (399)	3.0 (1.6-5.2), 12 (401)

Study Populations, Geographical Location	Active syphilis % Positive (95% CI), number positive (total number tested)						
	1998-1999 Round I	1999-2000 Round II	2000-2001 Round III	2002 Round IV	2003-2004 Round V	2004-2005 Round VI	2006 Round VII
Combined Injection Drug Users and Heroin Smokers: Southwest-A	ND	ND	ND	ND	ND	ND	1.0 (0.3-2.6), 4 (387)
Brothel Based Female Sex Workers: Central-B	13.8 (10.5-17.6), 54 (392)	6.2 (4.1-9.0), 25 (402)	8.1 (5.6-11.2), 33 (407)	3.9 (2.3-6.3), 16 (406)	3.2 (1.7-5.4), 13 (404)	1.7 (0.7-3.6), 7 (401)	1.5 (0.6-3.2), 6 (400)
Central-C	ND	10.2 (7.2-14.1), 33 (322)	ND	9.2 (5.1-15.0), 14 (152)	10.7 (6.4-16.6), 17 (159)	10.7 (6.2-16.7), 16 (150)	4.7 (1.9-9.4), 7 (150)
Central-D	ND	ND	14.8 (11.4-18.8), 57 (384)	6.7 (4.5-9.6), 27 (402)	6.0 (3.9-8.8), 24 (401)	3.8 (2.1-6.2), 15 (397)	4.2 (2.5-6.7), 17 (401)
Central-E	28.7 (23.5-34.4), 81 (282)	ND	ND	ND	ND	ND	ND
Central-L	ND	ND	ND	ND	11.0 (6.3-17.5), 15 (136)	8.4 (4.7-13.7), 14 (166)	6.0 (2.9-10.7), 10 (168)
Central-N	ND	ND	ND	ND	8.2 (5.7-11.5), 31 (376)	3.8 (2.1-6.3), 14 (370)	2.7 (1.3-4.9), 10 (373)
Central-P	ND	ND	ND	ND	12.2 (8.0-17.5), 25 (205)	8.4 (4.9-13.3), 16 (190)	6.3 (3.5-10.4), 14 (222)
Southwest-A, C [‡]	ND	11.7 (8.5-15.5), 41 (351)	6.0 (3.7-9.1), 20 (335)	5.0 (2.6-8.5), 12 (241)	4.8 (2.6-7.9), 14 (293)	2.8 (1.1-5.6), 7 (252)	0.4 (0-2.1), 1 (260)
Southwest-B	ND	ND	8.0 (4.6-12.9), 15 (187)	3.6 (1.5-7.3), 7 (195)	7.6 (4.1-12.6), 13 (171)	4.2 (1.7-8.4), 7 (167)	2.9 (0.9-6.6), 5 (174)
South-E	ND	ND	ND	ND	5.1 (1.1-14.1), 3 (59)	9.7 (3.6-19.9), 6 (62)	1.9 (0-10.3), 1 (52)
Street Based Female Sex Workers: Central-A	33.8 (29.1-38.6), 135 (400)	24.3 (20.3-28.7), 103 (423)	16.7 (13.3-20.6), 70 (419)	8.4 (5.9-11.6), 34 (403)	9.7 (7.0-13.1), 39 (401)	6.2 (4.1-9.0), 25 (402)	7.0 (4.7-10.0), 27 (386)
Central-B	ND	ND	ND	(1.1-6.4), 6 (199)	ND	ND	ND
Southeast-A	ND	ND	ND	ND	11.9 (8.9-15.5), 48 (402)	7.5 (5.1-10.5), 30 (402)	10.1 (7.4-13.5), 41(405)
Southwest-A	ND	ND	ND	4.7 (2.7-7.7), 15 (317)	1.5 (0.5-3.2), 6 (403)	ND	ND
Hotel Based Female Sex Workers: Central-A	ND	ND	ND	4.9 (3.0-7.5), 20 (405)	4.5 (2.7-7.0), 18 (400)	ND	ND
Southeast-A	ND	ND	ND	ND	5.3 (2.2-10.6), 7 (132)	1.6 (0.2-5.5), 2 (128)	4.2 (1.4-9.6), 5 (118)
Northeast-A	ND	ND	ND	ND	5.4 (2.5-10.0), 9 (166)	6.1 (2.9-10.9), 10 (165)	8.3 (4.6-13.5), 14 (169)
Casual Female Sex Workers: Southeast-D	ND	ND	ND	ND	ND	9.3 (4.3-16.9), 9 (97)	5.7 (1.9-12.8), 5 (88)
Southeast-H1	ND	ND	ND	ND	ND	10.0 (5.7-16.0), 15 (150)	3.5 (1.4-7.1), 7 (200)
Northwest-K1	ND	ND	ND	ND	6.9 (2.8-13.8), 7 (101)	4.2 (1.4-9.5), 5 (120)	4.7 (1.7-9.9), 6 (128)
Northwest-M1	ND	ND	ND	ND	1.0 (0.3-2.7), 4 (381)	0.5 (0-2.8), 1 (200)	0.9 (0.1-3.0), 2 (235)
South-A	ND	ND	ND	ND	5.1 (2.5-9.1), 10 (197)	1.5 (0.6-3.2), 6 (400)	1.5 (0.6-3.3), 6 (397)

Study Populations, Geographical Location	Active syphilis % Positive (95% CI), number positive (total number tested)						
	1998-1999 Round I	1999-2000 Round II	2000-2001 Round III	2002 Round IV	2003-2004 Round V	2004-2005 Round VI	2006 Round VII
Male Sex Workers (MSW): Central-A	ND	ND	7.7 (5.0-11.3), 24 (310)	3.2 (1.7-5.5), 13 (401)	6.2 (3.7-9.7), 17 (274)	3.8 (1.8-7.1), 9 (235)	4.9 (2.7-8.1), 14 (284)
Males Who Have Sex With Males (MSM): Central-A	ND	ND	1.8 (0.7-3.6), 7 (399)	0.7 (0.2-2.1), 3 (406)	1.5 (0.6-3.2), 6 (399)	2.0 (0.9-3.9), 8 (405)	0.2 (0-1.4), 1 (401)
MSM and MSW combined [§] : Central-A	7.0 (4.7-9.9), 28 (401)	6.7 (4.4-9.7), 26 (388)	ND	ND	ND	ND	ND
Central-C	ND	ND	ND	2.3 (1.0-4.2), 9 (400)	2.5 (1.2-4.5), 10 (400)	ND	ND
Southeast-A	ND	ND	ND	4.3 (2.5-6.8), 17 (397)	2.8 (1.4-4.9), 11 (398)	4.9 (2.7-8.2), 14 (283)	ND
Northeast-A	ND	ND	ND	3.0 (1.6-5.2), 12 (402)	3.3 (1.7-5.5), 13 (400)	5.6 (3.0-9.4), 13 (231)	ND
Hijras: Central-A	ND	ND	ND	10.4 (7.6-13.9), 41(393)	ND	ND	ND
Central-A, G	ND	ND	ND	ND	10.4 (7.6-13.8), 42 (405)	5.2 (1.1-3.2), 20 (381)	6.5 (4.2-9.6), 23 (353)
Partners of Hijra: Central-A, G	ND	ND	ND	ND	2.3 (0.3-8.0), 2 (88)	ND	ND
Babus (Brothel): Central-B	ND	ND	ND	1.6 (0.4-4.0), 4 (252)	2.0 (0.6-4.6), 5 (251)	ND	ND
Central-D	ND	ND	ND	6.0 (3.1-10.2), 12 (200)	6.3 (3.2-11.0), 11 (175)	ND	ND
Central-L	ND	ND	ND	ND	5.4 (1.1-14.9), 3 (56)	ND	ND
STI Patients: Central-A	11.0 (8.1-14.5), 44 (399)	5.2 (3.2-7.8), 21 (404)	ND	ND	ND	ND	ND
Southeast-A	7.6 (5.2-10.6), 31 (409)	4.2 (2.5-6.7), 17 (404)	2.2 (1.0-4.2), 9 (403)	ND	ND	ND	ND
Northwest-A, C*	2.2 (1.0-4.2), 9 (401)	1.7 (0.7-3.5), 7 (408)	1.5 (0.6-3.3), 6 (392)	ND	ND	ND	ND
Northeast-A	8.1 (5.6-11.2), 32 (397)	ND	5.1 (3.2-7.8), 20 (389)	0.9 (0-5.1), 1 (106)	ND	ND	ND
Truckers: Central-A	2.0 (0.9-3.9), 8 (403)	ND	2.1 (0.9-3.9), 9 (437)	1.1 (0.3- 2.5), 4 (402)	ND	ND	ND
Southwest-B	ND	ND	1.8 (0.7-3.6), 7 (392)	ND	ND	ND	ND
Southwest-B1	ND	ND	ND	ND	ND	0 (398)	ND
Dockworkers: Southeast-A	ND	ND	2.8 (1.4-5.0), 11 (392)	ND	ND	1.8 (0.7-3.6), 7 (395)	ND
Southwest-C	ND	ND	1.0 (0.3-2.5), 4 (401)	ND	ND	ND	ND

Study Populations, Geographical Location	Active syphilis % Positive (95% CI), number positive (total number tested)						
	1998-1999 Round I	1999-2000 Round II	2000-2001 Round III	2002 Round IV	2003-2004 Round V	2004-2005 Round VI	2006 Round VII
Rickshaw pullers: Central-A	ND	ND	ND	ND	0.2 (0-1.4), 1 (401)	0 (401)	ND
Southeast-A	ND	ND	1.0 (0.3-2.5), 4 (400)	ND	1.2 (0.4-2.9), 5 (401)	ND	ND
Southwest-B	ND	ND	1.0 (0.3-2.5), 4 (401)	ND	ND	ND	ND
Launch Workers: Central-A	ND	ND	ND	1.5 (0.5-3.2), 6 (402)	ND	ND	ND
TOTAL	11.3 (10.3-12.4), 440 (3886)	8.0 (7.2-8.8), 347 (4338)	4.6 (4.1-5.1), 322 (7063)	3.9 (3.5-4.4), 311 (7877)	4.5 (4.1-4.9), 471 (10445)	3.4 (3.1-3.8), 376 (11,029)	3.1 (2.8-3.5), 325 (10361)

†Central-A represents the merged result of Central-A1 and Central-A2

*Southwest A and C, two geographical related areas together representing one site

§In some sites male sex workers (MSW) and non-sex worker MSM could not be differentiated and they were sampled as a single group

*In the first round, sampling was done only in Northwest-A, in the subsequent rounds sampling was done for Northwest-A and C and these together represented a single site

ANNEXE: 2

Prevalence of HIV over six rounds of serological surveillance, 1998-1999, 1999-2000, 2000-2001, 2002, 2003-2004, 2004-2005 and 2006

Study Populations, Geographical Location	HIV % Positive (95% CI), number positive (total number tested)						
	1998-1999 Round I	1999-2000 Round II	2000-2001 Round III	2002 Round IV	2003-2004 Round V	2004-2005 Round VI	2006 Round VII
Injection Drug Users (IDU):							
Detoxification clinic: Central-A	2.5 (1.2-4.5), 10 (402)	0.2 (0-1.4), 1 (402)	0 (92)	ND	ND	ND	ND
Out of detoxification clinic: Central-A†	ND	1.4 (0.5-3.1), 6 (418)	1.7 (0.7-3.6), 7 (401)	4.0 (2.3-6.4), 16 (403)	4.0 (2.3-6.3), 16 (404)	4.9 (3.7-6.4), 52 (1061)	7.0 (5.5-8.7), 75 (1072)
Central-A1	ND	ND	ND	ND	ND	7.1 (5.2-9.3), 47 (664)	10.5 (8.3-13.1), 71 (674)
Central-A2	ND	ND	ND	ND	ND	1.3 (0.4-2.9), 5 (397)	1.0 (0.3-2.3), 4 (398)
Central-C	ND	ND	ND	ND	ND	0 (395)	0, (301)
Central-E	ND	ND	ND	ND	0 (107)	0 (103)	1.0 (0-5.2), 1 (105)
Central-H	ND	ND	ND	ND	0 (122)	0 (178)	0 (160)
Southeast-D	ND	ND	ND	ND	0 (86)	0.6 (0-3.5), 1 (159)	1.1 (0.1-4.0), 2 (178)
Southeast-H1	ND	ND	ND	ND	ND	0 (155)	0 (120)
Northwest-A	ND	0 (416)	0 (402)	0 (405)	0 (394)	0 (398)	0 (393)
Northwest-B	ND	ND	0 (120)	0 (200)	0 (239)	0 (208)	0 (200)
Northwest-B1	ND	ND	ND	ND	0 (78)	ND	ND
Northwest-B2	ND	ND	ND	ND	0 (47)	0 (66)	0 (69)
Northwest-C	ND	ND	ND	ND	ND	ND	0 (187)
Northwest-D	ND	ND	ND	ND	ND	0 (120)	0 (193)
Northwest-F	ND	ND	ND	ND	0 (85)	0 (57)	0 (69)
Northwest-F1	ND	ND	ND	ND	0 (57)	2.0 (0.1-10.9) 1 (49)	1.8 (0-9.7), 1 (55)
Northwest-G	ND	ND	ND	ND	ND	0 (111)	0 (122)
Northwest-K	ND	ND	ND	ND	ND	ND	0 (279)
Southwest-B	ND	ND	ND	ND	ND	0 (100)	0 (132)
Southwest-D	ND	ND	ND	ND	ND	0 (201)	0 (226)
South-A	ND	ND	ND	ND	ND	0 (202)	0 (234)
Female IDU: Central-A, E, H	ND	ND	ND	ND	ND	0 (119)	0.8 (0-4.5), 1 (121)
Heroin Smokers: Central-A	ND	ND	ND	0 (387)	0.8 (0.2-2.2), 3 (391)	0.5 (0.1-1.8), 2 (399)	0 (401)

Study Populations, Geographical Location	HIV % Positive (95% CI), number positive (total number tested)						
	1998-1999 Round I	1999-2000 Round II	2000-2001 Round III	2002 Round IV	2003-2004 Round V	2004-2005 Round VI	2006 Round VII
Combined Injection Drug Users and Heroin Smokers: Southwest-A	ND	ND	ND	ND	ND	ND	0 (387)
Brothel Based Female Sex Workers: Central-B	0 (392)	0 (402)	0.5 (0.1-1.8), 2 (407)	0.2 (0-1.4), 1 (406)	0.5 (0.1-1.8), 2 (404)	0.2 (0-1.4), 1 (401)	0.3 (0-1.4), 1 (400)
Central-C	ND	0 (322)	ND	0 (152)	0 (159)	0.7 (0-3.7), 1 (150)	0.7 (0-3.7), 1 (150)
Central-D	ND	ND	0.3 (0-1.4), 1 (384)	0.7 (0-2.2), 3 (402)	0.5 (0.1-1.8), 2 (401)	0.3 (0-1.4), 1 (397)	0.2 (0-1.4), 1 (401)
Central-E	1.5 (0.4-3.8), 4 (267)	ND	ND	ND	ND	ND	ND
Central-L	ND	ND	ND	ND	0 (136)	0 (166)	0 (168)
Central-N	ND	ND	ND	ND	0 (376)	0 (370)	0 (373)
Central-P	ND	ND	ND	ND	0.5 (0-2.7), 1 (205)	0 (190)	0.5 (0-2.5), 1 (222)
Southwest-A, C [‡]	ND	0 (351)	0 (335)	0 (241)	0 (293)	0.4 (0-2.2), 1 (252)	0 (260)
Southwest-B	ND	ND	0.5 (0-2.9), 1 (187)	0.5 (0-2.8), 1 (195)	0.6 (0-3.2), 1 (171)	0.6 (0-3.3), 1 (167)	0 (174)
South-E	ND	ND	ND	ND	0 (59)	0 (62)	0 (52)
Street Based Female Sex Workers: Central-A	0 (400)	0.2 (0-1.3), 1 (423)	0.5 (0.1-1.7), 2 (419)	0.2 (0-1.4), 1 (403)	0.2 (0-1.4), 1 (401)	0.2 (0-1.4), 1 (402)	0.3 (0-1.4), 1 (386)
Central-B	ND	ND	ND	0 (199)	ND	ND	ND
Southeast-A	ND	ND	ND	ND	0 (402)	0 (402)	0 (405)
Southwest-A	ND	ND	ND	0 (317)	0 (403)	ND	ND
Hotel Based Female Sex Workers: Central-A	ND	ND	ND	0.2 (0-1.4), 1 (405)	0 (400)	ND	ND
Southeast-A	ND	ND	ND	ND	1.5 (0.2-5.4), 2 (132)	0 (128)	0 (118)
Northeast-A	ND	ND	ND	ND	0.6 (0-3.3), 1 (166)	0.6 (0-3.3), 1 (165)	0 (169)
Casual Female Sex Workers: Southeast-D	ND	ND	ND	ND	ND	0 (97)	0 (88)
Southeast-H1	ND	ND	ND	ND	ND	0 (150)	0 (200)
Northwest-K1	ND	ND	ND	ND	2.0 (0.2-7.0), 2 (101)	1.7 (0.2-5.9), 2 (120)	0.8 (0-4.3), 1 (128)
Northwest-M1	ND	ND	ND	ND	0 (381)	0 (200)	0 (235)
South-A	ND	ND	ND	ND	0 (197)	0 (400)	0 (397)

Study Populations, Geographical Location	HIV % Positive (95% CI), number positive (total number tested)						
	1998-1999 Round I	1999-2000 Round II	2000-2001 Round III	2002 Round IV	2003-2004 Round V	2004-2005 Round VI	2006 Round VII
Male Sex Workers (MSW): Central-A	ND	ND	0 (310)	0 (401)	0 (274)	0 (235)	0.7 (0.1-2.5), 2 (284)
Males Who Have Sex With Males (MSM): Central-A	ND	ND	0 (399)	0.2 (0-1.4), 1 (406)	0 (399)	0 (405)	0.2 (0-1.4), 1 (401)
MSM and MSW combined [§] : Central-A	0.2 (0-1.4), 1 (401)	0 (388)	ND	ND	ND	ND	ND
Central-C	ND	ND	ND	0 (400)	0 (400)	ND	ND
Southeast-A	ND	ND	ND	0 (397)	0.3 (0-1.4), 1 (398)	0.4 (0-2.0), 1 (283)	ND
Northeast-A	ND	ND	ND	0 (402)	0.3 (0-1.4), 1 (400)	0.4 (0-2.4), 1 (231)	ND
Hijras: Central-A	ND	ND	ND	0.8 (0.2-2.2), 3 (393)	ND	ND	ND
Central-A, G	ND	ND	ND	ND	0.2 (0- 1.4), 1 (405)	0.8 (0.2-2.3), 3 (381)	0.6 (0.1-2.0), 2 (353)
Partners of Hijra: Central-A, G	ND	ND	ND	ND	0 (88)	ND	ND
Babus (Brothel): Central-B	ND	ND	ND	0 (252)	0 (251)	ND	ND
Central-D	ND	ND	ND	0 (200)	0 (175)	ND	ND
Central-L	ND	ND	ND	ND	0 (56)	ND	ND
STI Patients: Central-A	0.3 (0-1.4), 1 (399)	0 (404)	ND	ND	ND	ND	ND
Southeast-A	0.2 (0-1.4), 1 (409)	0 (404)	0.2 (0-1.4), 1 (403)	ND	ND	ND	ND
Northwest-A, C*	0 (401)	0 (408)	0 (392)	ND	ND	ND	ND
Northeast-A	0 (397)	ND	0 (389)	0 (106)	ND	ND	ND
Truckers: Central-A	0 (403)	ND	0 (437)	0 (402)	ND	ND	ND
Southwest-B	ND	ND	0 (392)	ND	ND	ND	ND
Southwest-B1	ND	ND	ND	ND	ND	0 (398)	ND
Dockworkers: Southeast-A	ND	ND	0 (392)	ND	ND	0 (395)	ND
Southwest-C	ND	ND	0 (401)	ND	ND	ND	ND

Study Populations, Geographical Location	HIV % Positive (95% CI), number positive (total number tested)						
	1998-1999 Round I	1999-2000 Round II	2000-2001 Round III	2002 Round IV	2003-2004 Round V	2004-2005 Round VI	2006 Round VII
Rickshaw pullers: Central-A	ND	ND	ND	ND	0.2 (0-1.4), 1 (401)	0 (401)	ND
Southeast-A	ND	ND	0 (400)	ND	0 (401)	ND	ND
Southwest-B	ND	ND	0 (401)	ND	ND	ND	ND
Launch Workers: Central-A	ND	ND	ND	0 (402)	ND	ND	ND
TOTAL	0.4 (0.3-0.7), 17 (3871)	0.2 (0.1-0.4), 8 (4338)	0.2 (0.1-0.3), 14 (7063)	0.3 (0.2-0.5), 27 (7877)	0.3 (0.2-0.5), 35 (10445)	0.6 (0.5-0.8), 70 (11,029)	0.9 (0.7-1.1), 91 (10,368)

†Central-A represents the merged result of Central-A1 and Central-A2

*Southwest A and C, two geographical related areas together representing one site

§In some sites male sex workers (MSW) and non-sex worker MSM could not be differentiated and they were sampled as a single group

*In the first round, sampling was done only in Northwest-A, in the subsequent rounds sampling was done for Northwest-A and C and these together represented a single site